



## **Cambridge International Examinations**

Cambridge International General Certificate of Secondary Education

CANDIDATE NAME						
CENTRE NUMBER		CANDIDATE NUMBER				
CHEMISTRY			0620/32			
Paper 3 (Exten	ded)	October/November 2015				
			1 hour 15 minutes			
Candidates ans	swer on the Question Paper.					
No Additional M	Naterials are required.					

## **READ THESE INSTRUCTIONS FIRST**

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use an HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid.

DO **NOT** WRITE IN ANY BARCODES.

Answer all questions.

Electronic calculators may be used.

A copy of the Periodic Table is printed on page 12.

You may lose marks if you do not show your working or if you do not use appropriate units.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [ ] at the end of each question or part question.

The syllabus is approved for use in England, Wales and Northern Ireland as a Cambridge International Level 1/Level 2 Certificate.

This document consists of **12** printed pages.



Use your copy of the Periodic Table to help you answer some of these questions	
(a) Predict the formulae of the following compounds.	
(i) nitrogen fluoride	
(ii) phosphorus sulfide	
	[2]
(b) Deduce the formulae of the following ions.	
(i) selenide	
(ii) gallium	
	[2]
(c) Use the following ions to determine the formulae of the compounds.	
ions OH- Cr <sup>3+</sup> Ba <sup>2+</sup> SO <sub>4</sub> <sup>2-</sup>	
compounds	
(i) chromium(III) sulfate	
(ii) barium hydroxide	
	[2]
	[Total: 6]

2	• •	lluted air contains two oxides of carbon and two oxides of nitrogen. A major source of these lutants is motor vehicles.
	(i)	Describe how carbon dioxide and carbon monoxide are formed in motor vehicle engines.
	(::)	[3]
	(ii)	State <b>one</b> adverse effect of each of these gases.  [2]
	(iii)	Nitrogen monoxide, NO, is released by motor vehicle exhausts.  Explain how nitrogen monoxide is formed in motor vehicle engines.
	(iv)	
		Suggest an explanation why this happens.  [1]
		edict the possible adverse effect on the environment when this non-metal oxide, $NO_2$ , reacts h water and oxygen.
		[2]
	` '	w are the amounts of carbon monoxide and nitrogen monoxide emitted by modern motor nicles reduced? Include an equation in your answer.
		[Total: 13]

**3** Two of the main uses of zinc are for galvanising and for making alloys.

One of the main ores of zinc is zinc blende, ZnS. There are two stages in the extraction of zinc from this ore.

(a)	Stage 1 Zinc oxide is made from zinc blende.
	Describe how this is done and write a word equation for the reaction.
	[2
(b)	Stage 2 Zinc oxide is reduced to zinc.
	Write a word equation for the reduction of zinc oxide by coke.
	[1
(c)	The zinc produced by this process is impure. It can be purified by electrolysis using a method which is similar to the purification of copper. Under the conditions used in the process, zinc is the product at the negative electrode (cathode).
	Complete the following description of this purification.
	The electrolyte is aqueous [1
	The negative electrode (cathode) is made of
	The positive electrode (anode) is impure zinc.
	The equation for the reaction at the cathode is
	The equation for the reaction at the anode is
	Explain why the concentration of the electrolyte does <b>not</b> change.
	[2

(d)	Bra	ass is an alloy which contains zinc.									
	(i)	Name the other metal in brass.									
	(ii)	Suggest <b>two</b> reasons why an alloy such as brass is preferred to either of its constituent metals.									
		[2]									
(e)	pie	an experiment to investigate the rate of rusting of steel, three pieces of steel were used. One ce of steel was completely coated with copper, one piece completely coated with zinc and third piece was left uncoated. All three pieces were left exposed to the atmosphere.									
	(i)	Explain why the uncoated piece started to rust.									
		[1]									
	(ii)	The coating on both of the other two pieces was scratched, exposing the steel.									
		exposed steel thin layer does not rust of zinc									
		steel									
		The piece of steel coated with zinc still did not rust but the copper-coated piece of steel rusted very rapidly.									
		Explain these observations in terms of the formation of ions and the transfer of electrons.									
		[4]									
		[Total: 17]									

4	(a)	Propane	reacts	with	chlorine	to	form	a	mixture	of	chloropropanes.	This i	s a	photochemica
		reaction.												

(i)	What is meant by the phrase photochemical reaction?
	[1

(ii) The products of this reaction include two isomers, one of which has the following structural formula.

Draw the structural formula of the other isomer.

		[1]
(iii)	Explain why these two different compounds are isomers.	
		[2]

**(b)** Bond breaking is an endothermic change and bond forming is an exothermic change.

Bond energy is the amount of energy in kJ/mol needed to break one mole of the specified bond.

Use the following bond energies to determine whether this reaction is exothermic or endothermic. You must show your reasoning.

bond	bond energies in kJ/mol
C-C1	338
C–H	412
Cl-Cl	242
H–C1	431
C–C	348

| <br> |
|------|------|------|------|------|------|------|------|------|
|      |      |      |      |      |      |      |      |      |
|      |      |      |      |      |      |      |      |      |
|      |      |      |      |      |      |      |      |      |
|      |      |      |      |      |      |      |      |      |
| <br> |
| <br> |
								FO:
								1:31

(c)	(i)	Chloropropane can be hydrolysed to propanol, $\mathrm{CH_3CH_2CH_2OH}$ , by sodium hydroxide.	
		Write the equation for this reaction.	
			[2]
	(ii)	Propanol can be dehydrated. It loses a water molecule to form a hydrocarbon.	
		Give the name and structural formula of this hydrocarbon.	
		name	
		structural formula	
			[2]
(	(iii)	Propanol is oxidised to a carboxylic acid by acidified potassium manganate(VII).	
		Deduce the name of this acid.	
			[1]
(d)	Pro	panol reacts with methanoic acid to form the ester propyl methanoate.	
		$CH_3CH_2CH_2OH + HCOOH \rightarrow HCOOCH_2CH_2CH_3 + H_2O$	
	4.0	g of methanoic acid was reacted with 6.0 g of propanol.	
	(i)	Calculate the $M_r$ of methanoic acid =	[1]
	(ii)	Calculate the $M_r$ of propanol =	[1]
(	(iii)	Determine which one is the limiting reagent. Show your reasoning.	
	/iv/\	Calculate the maximum yield in grams of propyl methanoate, $M_r$ = 88.	[—]
	(iv)	Calculate the maximum yield in grains of propyr methanoate, M <sub>r</sub> = 00.	[4]
			[1]

5 Iron is extracted from its ore, hematite, in a blast furnace.

Substances added to the furnace are:

- iron ore, hematite, containing impurities such as silica, SiO<sub>2</sub>
- air
- coke, C
- limestone, CaCO<sub>3</sub>

Substances formed in the blast furnace are:

- molten iron
- molten slag
- waste gases such as carbon dioxide

State the <b>two</b> functions of the coke used in the blast furnace.	
Write an equation for the conversion of hematite, Fe <sub>2</sub> O <sub>3</sub> , to iron.	
	[2]
Explain how the silica impurity is removed and separated from the molten iron.	
	[3]
The molten iron from the furnace is impure. t contains impurities which include the element carbon.	
Explain how the carbon is removed. Include an equation in your answer.	
	[3]
/	Vrite an equation for the conversion of hematite, Fe <sub>2</sub> O <sub>3</sub> , to iron.  Explain how the silica impurity is removed and separated from the molten iron.  The molten iron from the furnace is impure.  Contains impurities which include the element carbon.  Explain how the carbon is removed. Include an equation in your answer.

[Total: 10]

6 The table below shows the elements in the third period of the Periodic Table, the number of electrons in their outer energy level, their oxidation state in their common compounds and their melting points.

element	Na	Mg	Al	Si	Р	S	Cl	Ar
number of outer electrons	1	2	3	4	5	6	7	8
oxidation state	+1	+2	+3	+4/-4	<b>–</b> 3	-2	<b>–</b> 1	0
melting point/°C	98	650	660	1414	317	115	-101	-189

(a)	Describe and explain the variation in oxidation state across the period.	
		[3]
(b)	The first three elements, Na, Mg and A <i>l</i> , are metals.	
	Describe the structure of a typical metal.	
		[3]
(c)	Explain why Na, Mg and A $\it l$ are good conductors of electricity.	
		[1]
(d)	Which element exists as diatomic molecules of the type X <sub>2</sub> ?	
(α)		[1]
(e)	Silicon has a similar structure to diamond.	
	Explain why silicon has the highest melting point in the period.	
		[2]

(f)	Sodium chloride is a crystalline solid with a high melting point. It dissolves in water to give a neutral solution. Phosphorus trichloride is a liquid at room temperature. It reacts with water to form an acidic solution.
	Suggest an explanation for these differences in properties.
	[2]
(g)	Describe how you could show that magnesium oxide is a basic oxide and not an amphoteric oxide.
	[2]
(h)	Draw a dot-and-cross diagram showing the bonding in magnesium oxide. Show outer electrons only.

[3]

[Total: 17]

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge International Examinations Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cie.org.uk after the live examination series.

## DATA SHEET The Periodic Table of the Elements

							1110 1 0	10010 10	able of th	10 =10111							
								Gr	oup								
I	II											III	IV	V	VI	VII	0
							1 H Hydrogen										4 He Helium
7 <b>Li</b> Lithium	9 <b>Be</b> Beryllium											11 <b>B</b> Boron 5	12 <b>C</b> Carbon 6	14 N Nitrogen	16 O Oxygen 8	19 <b>F</b> Fluorine	20 <b>Ne</b> Neon
23 <b>Na</b> Sodium	Mg Magnesium 12											27 <b>A1</b> Aluminium 13	28 <b>Si</b> Silicon	31 P Phosphorus 15	32 <b>S</b> Sulfur 16	35.5 <b>C1</b> Chlorine 17	40 <b>Ar</b> Argon
39 <b>K</b> Potassium 19	40 <b>Ca</b> Calcium 20	45 Sc Scandium 21	48 <b>Ti</b> Titanium 22	51 <b>V</b> Vanadium 23	52 Cr Chromium 24	55 Mn Manganese 25	56 <b>Fe</b> Iron	59 Co Cobalt 27	59 <b>Ni</b> Nickel 28	64 Cu Copper 29	65 <b>Zn</b> Zinc 30	70 <b>Ga</b> Gallium 31	73 <b>Ge</b> Germanium 32	75 As Arsenic 33	79 <b>Se</b> Selenium 34	Bromine 35	84 <b>Kr</b> Krypton 36
85 <b>Rb</b> Rubidium	88 <b>Sr</b> Strontium 38	89 Y Yttrium 39	91 <b>Zr</b> Zirconium 40	93 <b>Nb</b> Niobium	96 <b>Mo</b> Molybdenum 42	Tc Technetium 43	101 <b>Ru</b> Ruthenium 44	103 <b>Rh</b> Rhodium 45	106 Pd Palladium 46	108 <b>Ag</b> Silver 47	112 Cd Cadmium 48	115 In Indium 49	119 <b>Sn</b> Tin 50	122 Sb Antimony 51	128 <b>Te</b> Tellurium 52	127 <b>I</b> lodine 53	131 <b>Xe</b> Xenon
133 Cs Caesium 55	137 <b>Ba</b> Barium	139 <b>La</b> Lanthanum  57 *	178 <b>Hf</b> Hafnium 72	181 <b>Ta</b> Tantalum 73	184 <b>W</b> Tungsten 74	186 <b>Re</b> Rhenium 75	190 Os Osmium 76	192 <b>Ir</b> Iridium	195 Pt Platinum 78	197 <b>Au</b> Gold 79	201 <b>Hg</b> Mercury	204 <b>T <i>l</i></b> Thallium 81	207 <b>Pb</b> Lead 82	209 <b>Bi</b> Bismuth	Po Polonium 84	At Astatine 85	Rn Radon 86
Fr Francium 87	226 <b>Ra</b> Radium 88	227 Ac Actinium 89 †															
*58-71 Lanthanoid series  †90-103 Actinoid series  Ce Cerium 58  140 Pr Neodymium 59  Neodymium 59  Neodymium 60			Nd Neodymium	Pm Promethium 61	150 <b>Sm</b> Samarium 62	152 <b>Eu</b> Europium 63	157 <b>Gd</b> Gadolinium 64	159 <b>Tb</b> Terbium 65	162 <b>Dy</b> Dysprosium 66	165 <b>Ho</b> Holmium 67	167 <b>Er</b> Erbium 68	169 <b>Tm</b> Thulium 69	173 <b>Yb</b> Ytterbium 70	175 <b>Lu</b> Lutetium 71			
Key a a = relative atomic mass X = atomic symbol b = proton (atomic) number			bol	232 <b>Th</b> Thorium 90	Pa Protactinium 91	238 <b>U</b> Uranium 92	Np Neptunium 93	Pu Plutonium 94	Am Americium 95	Cm Curium 96	<b>Bk</b> Berkelium  97	Cf Californium 98	Es Einsteinium 99	Fm Fermium 100	Md Mendelevium 101	No Nobelium 102	Lr Lawrencium 103

The volume of one mole of any gas is 24 dm<sup>3</sup> at room temperature and pressure (r.t.p.).