



Cambridge International Examinations

Cambridge International General Certificate of Secondary Education

| CHEMISTRY | | | 062 | 20/32 |
|-------------------|--|---------------------|-----|-------|
| CENTRE NUMBER | | CANDIDATE NUMBER | | |
| CANDIDATE NAME | | | | |

Paper 3 (Extended)

May/June 2014

1 hour 15 minutes

Candidates answer on the Question Paper.

No Additional Materials are required.

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use an HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid.

DO NOT WRITE IN ANY BARCODES.

Answer all questions.

Electronic calculators may be used.

A copy of the Periodic Table is printed on page 12.

You may lose marks if you do not show your working or if you do not use appropriate units.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [] at the end of each question or part question.

The syllabus is approved for use in England, Wales and Northern Ireland as a Cambridge International Level 1/Level 2 Certificate.



1 The table below gives the electron distributions of atoms of different elements.

| element | electron distribution |
|---------|-----------------------|
| Α | 2 + 7 |
| В | 2 + 8 + 4 |
| С | 2+8+8+1 |
| D | 2 + 8 + 18 + 5 |
| E | 2 + 8 + 18 + 7 |
| F | 2 + 8 + 18 + 18 + 8 |

For each of the following, select an element or elements from the table that matches the description. Each element may be selected once, more than once or not at all.

| (a) | These two elements are in the same group. | |
|-----|--|-----|
| | | [1] |
| (b) | This element forms a fluoride with a formula of the type XF ₃ . | |
| | | [1] |
| (c) | This element reacts violently with cold water. | |
| | | [1] |
| (d) | This element has a macromolecular structure similar to that of diamond. | |
| | | [1] |
| (e) | The only oxidation state of this element is 0. | |
| | | [1] |
| (f) | This element is bromine. | |
| | | [1] |
| (g) | This element is a good conductor of electricity. | |
| | | [1] |

[Total: 7]

2

| (a) Nat | tural gas, which is mainly methane, is a fossil fuel. |
|------------------|---|
| (i) | What is meant by the term <i>fuel</i> ? |
| | [1] |
| (ii) | Name two other fossil fuels. |
| | [2] |
| (iii) | Name a solid fuel which is not a fossil fuel. |
| | [1] |
| | ssil fuels are formed by the anaerobic decomposition of organic matter. Anaerobic means in absence of oxygen. |
| (i) | The organic matter contains hydrogen and carbon. Suggest the products that would be formed if the decomposition occurred in the presence of oxygen. |
| | [2] |
| (ii) | What are the two main disadvantages in the widespread use of fossil fuels? |
| | |
| | |
| | [2] |
| | [Total: 8] |

| 3 | | Plant growth is improved by the availability of essential elements, such as nitrogen, and by the soil naving a suitable pH. | | | | | | |
|---|------|---|---|------------|--|--|--|--|
| | (a) | | ogen-based fertilisers are made from ammonia. Ammonia is manufactured by the Habecess. | er | | | | |
| | | (i) | Describe the Haber process giving reaction conditions and a balanced equation. (Do not discuss reaction rate and yield.) | | | | | |
| | | | | | | | | |
| | | | | | | | | |
| | | | | | | | | |
| | | | [5 | 5] | | | | |
| | | (ii) | Fertilisers contain nitrogen. Name the other two elements essential for plant growth commonly found in fertilisers. | | | | | |
| | (I-) | 0 | | 2] | | | | |
| | (D) | Cro | ps do not grow well if the soil is too acidic. | | | | | |
| | | (i) | One cause of acidity in soil is acid rain. Explain how acid rain is formed. | | | | | |
| | | | | | | | | |
| | | | | | | | | |
| | | (ii) | Name two bases which are used to increase the pH of acidic soils. | 3] | | | | |
| | | | [2 | 2] | | | | |

[Total: 12]

4

| Propanoic acid is a carboxylic acid. Its formula is CH ₃ –CH ₂ –COOH. | | | | | | | |
|---|------|--|-----|--|--|--|--|
| (a) | Pro | Propanoic acid is the third member of the homologous series of carboxylic acids. | | | | | |
| | (i) | Give the name and structural formula of the fourth member of this series. | | | | | |
| | | name | | | | | |
| | | formula | [2] | | | | |
| | (ii) | Members of a homologous series have very similar chemical properties. State three other characteristics of a homologous series. | | | | | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | |
| | | | [3] | | | | |
| (b) | Car | boxylic acids can be made by the oxidation of alcohols. | | | | | |
| | (i) | Draw the structural formula of the alcohol which can be oxidised to propanoic acid. Show all atoms and bonds. | | | | | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | |
| | | | [1] | | | | |
| | (ii) | Name a reagent, other than oxygen, which can oxidise alcohols to carboxylic acids. | | | | | |
| | | | [2] | | | | |
| | | | | | | | |
| | | | | | | | |

(c) Complete the following equations for some of the reactions of propanoic acid.

(iii) LiOH + $CH_3CH_2COOH \rightarrow \dots + \dots + \dots$

| The | e salts of this acid are called propanoates. | |
|------|--|-----|
| (i) | zinc + propanoic acid \rightarrow + hydrogen | [1] |
| (ii) | calcium + propanoic \rightarrow + + | [1] |

[1]

[Total: 18]

(d) A piece of magnesium was added to 100 cm³ of an aqueous acid. The time taken for the metal to react completely was measured. This experiment was repeated using different aqueous acids. The same volume of acid was used in each experiment and the pieces of magnesium used were identical. In one experiment the reaction was carried out at a different temperature.

| experiment | acid | concentration in mol/dm³ | temperature /°C | time / minutes |
|------------|--------------|--------------------------|--------------------|-------------------|
| Α | propanoic | 1.0 | 20 | 5 |
| В | propanoic | 1.0 | 30 | 3 |
| С | propanoic | 0.5 | 20 | 8 |
| D | hydrochloric | 1.0 | 20 | 1 |

Explain the following in terms of collision rate between reacting particles.

| (i) | Why is the rate in experiment C slower than the rate in experiment A ? | |
|-------|--|-----|
| | | |
| | | |
| (ii) | Why is the rate in experiment B faster than the rate in experiment A ? | |
| | | |
| | | |
| (iii) | Why is the rate in experiment D faster than the rate in experiment A ? | |
| | | |
| | | [3] |

| 5 | Carbonyl | chloride i | s made | from ca | arbon r | monoxide | and chlorine | Э. |
|---|----------|------------|--------|---------|---------|----------|--------------|----|
| _ | | | | | | | | _ |

$$CO(g) + Cl_2(g) \rightleftharpoons COCl_2(g)$$

| (a) | | o methods of preparing carbon monoxide are from methane and oxygen, and from methane steam. |
|-----|-------|---|
| | (i) | The reaction between methane and oxygen can also form carbon dioxide. How can carbon monoxide be made instead of carbon dioxide? |
| | | [1] |
| | (ii) | The following reaction is used to make carbon monoxide and hydrogen. The reaction is carried out at 1100 °C and normal pressure. |
| | | $CH_4(g) + H_2O(g) \rightleftharpoons CO(g) + 3H_2(g)$ |
| | | The reaction is reversible and comes to equilibrium. Suggest why a high temperature is used. |
| | | |
| | | |
| | | [2] |
| | (iii) | What is the disadvantage of using a high pressure for the reaction given in (a)(ii)? |
| | | [2] |
| (b) | Des | orine is made by the electrolysis of concentrated aqueous sodium chloride. scribe this electrolysis. Write ionic equations for the reactions at the electrodes and name sodium compound formed. |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | [5] |

(c) The structural formula of carbonyl chloride is given below.



Draw a diagram showing the arrangement of the valency electrons around the atoms in one molecule of this covalent compound.

Use o to represent an electron from an oxygen atom.

Use x to represent an electron from a chlorine atom.

Use • to represent an electron from a carbon atom.

[3]

[Total: 13]


6 Scandium, proton number 21, is not a typical transition element.

| (a) | Scandium is a low density metal which has only one oxidation state in its compounds. Scandium compounds are white solids which form colourless solutions. Titanium, the next metal in the period, is a far more typical transition element. How would the properties of titanium differ from those of scandium? |
|-----|---|
| | |
| | |
| | |
| | [3] |

| (b) | Scandium fluoride is an ionic compound. The valency of scandium in scandium fluoride is three. |
|-----|---|
| | Draw a diagram which shows the formula of this compound, the charges on the ions and the arrangement of the valency electrons around the negative ions. |
| | Use × to represent an electron from a fluorine atom. Use o to represent an electron from a scandium atom. |
| | |

| | [3] |
|-----|---|
| (c) | Scandium oxide is insoluble in water. Describe how you could show that it is an amphoteric oxide. |
| | |
| | |
| | [3] |
| | [Total: 9] |

7 The soluble salt hydrated lithium sulfate is made by titration from the soluble base lithium hydroxide.



| (a) | The sulfuric acid is added slowly from the burette until the indicator just changes colour. The |
|-----|---|
| | volume of sulfuric acid needed to just neutralise the lithium hydroxide is noted. |
| | Describe how you would continue the experiment to obtain pure dry crystals of hydrated lithium |
| | sulfate. |

| | • |
|----|----|
| | - |
| aj | :1 |

(b) Using 25.0 cm³ of aqueous lithium hydroxide, concentration 2.48 mol/dm³, 2.20 g of hydrated lithium sulfate was obtained.

Calculate the percentage yield, giving your answer to one decimal place.

$$2 \text{LiOH + H}_2 \text{SO}_4 \ \rightarrow \ \text{Li}_2 \text{SO}_4 \ + \ 2 \text{H}_2 \text{O}$$

$$Li_2SO_4 + H_2O \rightarrow Li_2SO_4.H_2O$$

Number of moles of LiOH used =

Number of moles of Li₂SO₄.H₂O which could be formed =

Mass of one mole of Li_2SO_4 . $H_2O = 128g$

Maximum yield of Li_2SO_4 . $H_2O =g$

Percentage yield = [4]

(c) An experiment was carried out to show that the formula of the hydrated salt is Li₂SO₄·H₂O. A sample of the hydrated salt was weighed and its mass recorded. It was then heated and the

| the same. This procedure is called 'heating to constant mass'. | |
|---|------|
| (i) What is the reason for heating to constant mass? | (i) |
| [1] | |
| (ii) The mass of the hydrated salt is m ₁ and the mass of the anhydrous salt is m ₂ . Explain how you could show that the hydrated salt has one mole of water of crystallisation per mole of the anhydrous salt. | (ii) |
| | |
| [3] | |
| [Total: 13] | |

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included the publisher will be pleased to make amends at the earliest possible opportunity.

DATA SHEET The Periodic Table of the Elements

| Group | | | | | | | | | | | | | | | | | |
|-----------------------------------|-----------------------------|---|------------------------------------|-------------------------------------|-------------------------------------|-----------------------------------|------------------------------------|--------------------------------------|-----------------------------------|--------------------------------------|-----------------------------------|--|------------------------------------|-------------------------------------|-------------------------------------|-------------------------------------|----------------------------------|
| I | II | | | | | | | Oit | Бир | | | III | IV | V | VI | VII | 0 |
| 1 H Hydrogen | | | | | | | | | | | | | | | | 4 He Helium 2 | |
| 7 Li Lithium | 9 Be Beryllium | | | | | | | | | | | 11 B Boron 5 | 12 C Carbon | 14 N Nitrogen | 16 O Oxygen 8 | 19 F Fluorine | 20 Ne Neon |
| 23 Na Sodium | Mg Magnesium 12 | | | | | | | | | | | 27 A 1 Aluminium 13 | 28 Si Silicon | 31 P Phosphorus 15 | 32 S Sulfur | 35.5 C1 Chlorine 17 | 40 Ar Argon |
| 39 K Potassium 19 | 40 Ca Calcium 20 | 45 Sc Scandium 21 | 48 Ti Titanium 22 | 51 V Vanadium 23 | 52 Cr Chromium 24 | 55 Mn Manganese 25 | 56 Fe Iron | 59 Co Cobalt 27 | 59 Ni Nickel 28 | 64 Cu Copper 29 | 65 Zn Zinc | 70 Ga Gallium | 73 Ge Germanium 32 | 75 As Arsenic | 79 Se Selenium 34 | Br Bromine 35 | 84 Kr Krypton 36 |
| 85 Rb Rubidium 37 | 88 Sr Strontium 38 | 89 Y Yttrium | 91 Zr Zirconium 40 | 93 Nb Niobium 41 | 96 Mo Molybdenum 42 | Tc Technetium 43 | 101 Ru Ruthenium | 103 Rh Rhodium 45 | 106 Pd Palladium 46 | 108 Ag Silver | Cd Cadmium | 115 In Indium | 119 Sn Tin | 122 Sb Antimony 51 | 128 Te Tellurium 52 | 127 I lodine 53 | 131 Xe Xenon 54 |
| 133 Cs Caesium 55 | 137 Ba Barium | 139 La Lanthanum 57 * | 178 Hf Hafnium 72 | 181 Ta Tantalum 73 | 184 W Tungsten 74 | 186 Re Rhenium 75 | 190 Os Osmium 76 | 192 Ir Iridium | 195 Pt Platinum 78 | 197 Au Gold 79 | 201 Hg Mercury | 204 T <i>l</i> Thallium 81 | 207 Pb Lead | 209 Bi Bismuth | Po Polonium 84 | At Astatine 85 | Rn Radon 86 |
| Fr Francium 87 | 226 Ra Radium | 227 AC Actinium 89 † | | | | | | | | | | | | | | | |
| | | | 141 Pr Praseodymium 59 | 144 Nd Neodymium 60 | Pm Promethium 61 | 150 Sm Samarium 62 | 152 Eu Europium 63 | 157 Gd Gadolinium 64 | 159 Tb Terbium 65 | 162 Dy Dysprosium 66 | 165 Ho Holmium 67 | 167 Er Erbium 68 | 169 Tm Thulium 69 | 173 Yb Ytterbium 70 | 175 Lu Lutetium 71 | | |
| Key | X | = relative ator (= atomic sym = proton (aton | bol | 232 Th Thorium 90 | Pa Protactinium 91 | 238 U Uranium 92 | Np Neptunium 93 | Pu Plutonium 94 | Am Americium 95 | Cm Curium 96 | Bk Berkelium 97 | Cf Californium 98 | Es Einsteinium 99 | Fm Fermium 100 | Md Mendelevium 101 | No Nobelium 102 | Lr Lawrencium 103 |

The volume of one mole of any gas is 24 dm³ at room temperature and pressure (r.t.p.).